

---

# An Introduction To Ordinary Differential Equations

---

*An Introduction To  
Ordinary Differential  
Equations*

*Downloaded from  
[www.whlarchitecture.com](http://www.whlarchitecture.com)  
by Vazquez & Rory*

---

## UNDERSTANDING THE BASICS OF DIFFERENTIAL EQUATIONS

---

Mathematics often serves as the language through which we describe change. From the motion of planets to the growth of populations, change is governed by rates. An **ordinary differential equation** (ODE) is a mathematical equation that relates a

function of one independent variable to its derivatives. In simple terms, it expresses how a quantity changes over time or space. For students stepping into the world of calculus-based modeling, understanding what an ODE is and how to work with it forms a critical foundation for physics, engineering, economics, and biology.

The phrase *An Introduction To Ordinary Differential Equations* might sound intimidating, but at its core, the subject

is about predicting the future behavior of dynamic systems. By learning to set up and solve ODEs, you gain the ability to translate real-world phenomena into precise mathematical statements. This article will walk you through the essential concepts, classifications, solution methods, and practical significance of ODEs — all in a clear, step-by-step manner.

---

## WHAT IS AN ORDINARY DIFFERENTIAL EQUATION?

---

An ordinary differential equation is an equation that contains one or more derivatives of an unknown function  $y(x)$  with respect to a single independent variable  $x$ . The term **ordinary** distinguishes it from **partial differential equations**, which involve

derivatives with respect to multiple variables.

For example, consider the equation:

$$dy/dx = 3x^2$$

Here,  $y$  is the dependent variable,  $x$  is the independent variable, and  $dy/dx$  is the first derivative. Solving this ODE means finding a function  $y(x)$  that satisfies the relationship. In this case, integrating both sides gives  $y(x) = x^3 + C$ , where  $C$  is an arbitrary constant.

ODEs can also involve higher derivatives, such as second derivatives ( $d^2y/dx^2$ ) or even higher orders. The goal is always to determine the function that fits the equation given certain initial or boundary conditions.

# Key Components of an ODE

- **Independent variable:** Usually denoted by  $x$  or  $t$  (time).
- **Dependent variable:** Typically  $y$  or  $u$ , representing the quantity of interest.
- **Derivatives:** Indicate how quickly the dependent variable changes with respect to the independent variable.
- **Order:** The highest derivative present in the equation.
- **Linear vs. Nonlinear:** A classification based on whether the dependent variable and its derivatives appear linearly.

## ORDER AND DEGREE: THE FIRST CLASSIFICATIONS

The **order** of an ordinary differential equation is the order of the highest derivative. For instance,  $d^2y/dx^2 + 5 dy/dx + 6y = 0$  is a second-order ODE. The **degree** is the power of the highest derivative term after the equation has been cleared of radicals and fractions. Most introductory courses focus on first-order and second-order ODEs because they cover a vast range of practical problems.

## First-Order ODEs

A general first-order ODE can be written as  $dy/dx = f(x, y)$ . Examples include:

- **Separable equations:**  $dy/dx =$

$$g(x)h(y)$$

- **Linear equations:**  $dy/dx + P(x)y = Q(x)$
- **Exact equations:**  $M(x,y)dx + N(x,y)dy = 0$

Each type has its own standard solution technique, which we will explore below.

---

## WHY ARE ORDINARY DIFFERENTIAL EQUATIONS IMPORTANT?

---

ODEs appear everywhere in science and engineering. They model:

- **Population growth:**  $dP/dt = rP$  (exponential growth)
- **Radioactive decay:**  $dN/dt = -\lambda N$
- **Newton's law of cooling:**  $dT/dt = -k(T - T_{env})$

- **RC circuits:**  $dq/dt + (1/RC) q = V/R$
- **Mechanical oscillations:**  $m d^2x/dt^2 + c dx/dt + kx = F(t)$

Mastering the basics of ODEs empowers you to analyze how systems evolve, predict future states, and design control systems. In other words, an introduction to ordinary differential equations is a gateway to applied mathematics.

---

## SOLVING FIRST-ORDER ORDINARY DIFFERENTIAL EQUATIONS

---

Solution methods for first-order ODEs are often the first major topic in any introductory course. The choice of method depends on the form of the equation.

# 1. Separable Equations

An ODE is separable if it can be written in the form  $dy/dx = g(x)h(y)$ . The technique is to separate variables:

1. Rewrite as  $dy / h(y) = g(x) dx$ .
2. Integrate both sides.
3. Solve for  $y$  if possible.

**Example:** Solve  $dy/dx = 2xy$ .

Separate:  $dy/y = 2x dx$ . Integrate:  $\ln|y| = x^2 + C$ . Hence,  $y = Ce^{(x^2)}$ .

# 2. Linear First-Order ODEs

A linear first-order ODE has the standard form  $dy/dx + P(x)y = Q(x)$ . The solution uses an **integrating factor**  $\mu(x) = e^{(\int P(x)dx)}$ . Multiply both sides by  $\mu(x)$ , then the left side becomes  $d/dx(\mu y)$ . Integrate to find  $y$ .

**Example:** Solve  $dy/dx - (2/x)y = x^2$  for  $x > 0$ .

Here  $P(x) = -2/x$ , so  $\mu(x) = e^{(\int -2/x dx)} = e^{(-2 \ln x)} = x^{-2}$ . Multiply:  $x^{-2} dy/dx - 2x^{-3}y = 1$ . Left side is  $d/dx(x^{-2}y) = 1$ . Integrate:  $x^{-2}y = x + C$ , so  $y = x^3 + Cx^2$ .

## ORDINARY DIFFERENTIAL

# 3. Exact Equations

An equation  $M(x,y)dx + N(x,y)dy = 0$  is exact if  $\partial M/\partial y = \partial N/\partial x$ . The solution is a function  $F(x,y) = C$  such that  $dF = M dx + N dy$ . Finding  $F$  involves integrating  $M$  with respect to  $x$  and then comparing with  $N$ .

These three methods cover the most common first-order ODEs encountered in applications. More advanced techniques, such as substitution (e.g., Bernoulli equations) or numerical methods, build on these fundamentals.

---

## SECOND-ORDER LINEAR

## EQUATIONS

---

Second-order ODEs are crucial for modeling oscillations, wave propagation, and electrical circuits. A general second-order linear ODE is written as:

$$a(x) \frac{d^2y}{dx^2} + b(x) \frac{dy}{dx} + c(x) y = f(x)$$

If  $f(x) = 0$ , the equation is **homogeneous**; otherwise, it is **nonhomogeneous**. For constant-coefficient equations ( $a$ ,  $b$ ,  $c$  are constants), we can find solutions using characteristic equations.

## Homogeneous

## Case: $ay'' + by' + cy = 0$

Assume a solution of the form  $y = e^{(rx)}$ . Substituting leads to the characteristic equation  $ar^2 + br + c = 0$ . Depending on the discriminant, we get:

- **Two distinct real roots:**  $y = C_1 e^{(r_1 x)} + C_2 e^{(r_2 x)}$
- **Repeated real root:**  $y = (C_1 + C_2 x) e^{(r x)}$
- **Complex conjugate roots:**  $y = e^{(ax)}(C_1 \cos \beta x + C_2 \sin \beta x)$

## Nonhomogeneous Case

To solve  $ay'' + by' + cy = f(x)$ , we first find the complementary solution (from the homogeneous equation) and then a particular solution. Methods include **undetermined coefficients** (for certain simple  $f(x)$  like polynomials, exponentials, sines/cosines) and **variation of parameters** (more general).

**Example:** Solve  $y'' - 3y' + 2y = 2e^{(3x)}$ .

Homogeneous solution: characteristic  $r^2 - 3r + 2 = 0$  gives  $r=1,2$ , so  $y_h = C_1 e^x + C_2 e^{(2x)}$ . For particular solution, try

$y_p = A e^{(3x)}$ . Substituting gives  $9A - 9A + 2A = 2 \Rightarrow 2A = 2 \Rightarrow A=1$ . So  $y_p = e^{(3x)}$ . General solution:  $y = C_1 e^x + C_2 e^{(2x)} + e^{(3x)}$ .

---

## INITIAL VALUE PROBLEMS AND BOUNDARY VALUE PROBLEMS

---

An ODE alone yields a family of solutions. To pin down a specific solution, we need additional conditions:

- **Initial value problem (IVP):**

Conditions are specified at a single point, typically the starting time. For example,  $y(0) = 1$  and  $y'(0) = 0$ .

- **Boundary value problem**

**(BVP):** Conditions are given at two different points, often the endpoints of a spatial interval. For

example,  $y(0) = 0$  and  $y(L) = 0$ .

Most introductory courses focus on IVPs because they align with time evolution problems.

---

## PRACTICAL APPLICATIONS OF ODES

---

The true power of an introduction to ordinary differential equations lies in its applicability. Let's look at a few classic models:

# Population Dynamics

The logistic equation  $dP/dt = rP(1 - P/K)$  models population growth with limited resources. It is a first-order nonlinear



ODE, solvable by separation of variables. The solution yields an S-shaped curve that levels off at the carrying capacity  $K$ .

## Spring-Mass

## Systems

Consider a mass attached to a spring. By Newton's second law, the displacement  $x(t)$  satisfies  $m d^2x/dt^2 + c dx/dt + kx = F(t)$ . This second-order linear ODE explains harmonic motion, damping